

RIPC prior to cardiac surgery

Dr Derek J Hausenloy

BHF Senior Clinical Research Fellow
The Hatter Cardiovascular Institute,
University College London, UK.

Honorary Consultant Cardiologist
University College London Hospital, UK.

IC Therapeutics Meeting, Monday November 15th 2010.

- Novel cardioprotective strategies are required to improve clinical outcomes in CHD patients.
- Sicker patients are being operated on.
- CK-MB/Troponin release during CABG surgery linked to worse outcomes. Renal and cerebral outcomes.
- RIPC at proof-of-concept stage.
- Multicentre large randomized clinical trials needed.

'Conditioning' the heart *remotely*

Remote Ischemic Preconditioning



Heart Ischemia

Heart Reperfusion

No 'Conditioning'



'Conditioned'




CABG surgery
Cardiac transplant
PCI

STEMI

CABG surgery
Cardiac transplant
Cardiac arrest
STEMI

From heart to inter-organ protection

Hausenloy & Yellon Cardiovasc Res 2008;79;377.



RIPC in paediatric cardiac surgery

Cheung et al JACC 2006;47;2277.

- Complicated surgery with high morbidity.
- 37 children requiring corrective cardiac surgery.
- RIPC comprised inflation of cuff placed on lower limb to $15\text{mmHg} > \text{SBP}$ for 4x5 min.
- RIPC improved troponin-I, inotrope requirement and airway resistance.



RIPC in CABG surgery

Hausenloy et al Lancet 2007:370;575.



- 57 adult CABG patients:
RIPC- 3x5 min cuff inflation
Control- 30 min deflated cuff
- RIPC reduced myocardial injury by 43%.
- However, mixture of ICCF and cardioplegia.




Time (h)

RIPC in CABG patients (cardioplegia)

Venugopal et al Heart 2009 95(19):1567-71


- Cardioplegia only patients.
- 36 adult CABG patients
RIPC-3x5 min cuff inflation
Control-30 min deflated cuff
- RIPC reduced myocardial injury by 39% (reduction in trop T AUC over 72 hrs).



RIPC on renal outcomes

Venugopal et al Am J Kidney Dis. 2010 Oct 23. [Epub ahead of print]


- Acute kidney injury (AKI) affects up to 30% of patients undergoing cardiac surgery, with 1%-2% of patients going on to require dialysis therapy.
- 78 adult elective CABG patients randomised to:
RIPC-3x5 min cuff inflation Control-30 min deflated cuff
- RIPC reduced the development of AKI from 10/40 patients in control to 4/38 in RIPC-treated patients.



RIPC in diabetic CABG patients

Babu et al Unpublished

- The diabetic heart is resistant to IPC.
- 55 adult diabetic CABG patients RIPC vs control.
- RIPC had no effect on myocardial injury .



Summary of RIPC CABG studies



Study	N	RIC	Pt	Cardio protect	Clamp time	CPB time	Result	Notes
Cheung et al 2006	37	4x5 min leg	Children	Cardio	55,59	80,88	↓Trop I	
Hausenloy et al 2007	57	3x5 min arm	Adult CABG± valve	CCF and cardio	45,36	80,75	43%↓ 72 hr AUC Trop T	
Venugopal et al 2009	45	3x5 min arm	Adult CABG± Valve	Cardio	65,53	97,86	42%↓ 72 hr AUC Trop T	No diabetics
Gegouskov et al 2009 abstract	40	3x5 min arm	Adult CABG± valve	Cardio	N/A	N/A	↓6, 24, 48 hr CKMB and Trop T	
Thielmann et al 2010	53	3x5 min arm	Adult CABG	Cryst	76,71	110,109	35%↓ 48 hr AUC Trop I	No diabetics
Rahman et al 2010	162	3x5 min arm	Elective urgent CABG	Cardio	71, 76	96,100	No difference in 48hr Trop T	IV GTN given to most patients. No diabetics.
Wagner et al 2010	101	3x5 min arm 18 hr pre	Elective CABG	Cryst	51, 45	87,80	↓Trop I	



Effect of Remote Ischaemic Conditioning on clinical outcomes in patients undergoing Coronary Artery bypass graft surgery

Hypothesis: RIC improves clinical outcomes in patients (Euroscore>6) undergoing CABG surgery

Design: 11 centre 1610 patient randomised double-blinded RCT

Funding: £1.6 million NIHR/MRC/BHF- EME grant

Primary outcome: Death, MI, revascularisation, stroke at one year.

Secondary outcomes:

- Peri-operative myocardial injury (high-sens Trop-T)
- Peri-operative renal injury (AKI score, blood NGAL)
- Length of ITU/hospital stay and inotrope score
- SMWT, Quality of life, LV ejection fraction

Why the discrepancy?

Patient selection:

- Routine versus urgent
- CABG±valve
- Diabetics

RIC stimulus:

- Arm versus leg
- Number of cycles
- Applied effectively

Cardiac surgery:

- Isoflurane
- IV GTN
- Cross-clamp and bypass times

- Remote ischaemic preconditioning has emerged as a non-invasive virtually cost-free strategy for protecting the heart during CABG surgery- proof-of-concept.
- The optimal RIPC protocol and patient group remains to be determined.
- Large multi-centre clinical studies are required to determine whether any effect on clinical outcomes.

Acknowledgements

Professor Derek Yellon
Dr Vinod Venugopal
Dr Girish Babu
Dr Edney Boston-Griffiths

Staff and patients at the Heart
Hospital

British Heart Foundation

